

This is a repository copy of *The potential impact of human visceral leishmaniasis vaccines on population incidence*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/162741/>

Version: Accepted Version

Article:

Le Rutte, Epke, Coffeng, Luc, Malvoti, Stefano et al. (2 more authors) (Accepted: 2020)
The potential impact of human visceral leishmaniasis vaccines on population incidence.
PLOS NEGLECTED TROPICAL DISEASES. ISSN 1935-2735 (In Press)

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

The potential impact of human visceral leishmaniasis vaccines on population incidence

Epke A. Le Rutte^{1,2,3*}, Luc E. Coffeng¹, Stefano Malvoti⁴, Paul M. Kaye⁵, Sake J. de Vlas¹.

¹ Department of Public Health, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands

² Department of Epidemiology and Public Health, Swiss Tropical and Public Health Institute, Basel, Switzerland

³ University of Basel, Basel, Switzerland

⁴ MM Global Health Consulting, Zurich, Switzerland

⁵ York Biomedical Research Institute, Hull York Medical School, University of York, York, United Kingdom

*Corresponding author: e.lerutte@erasmusmc.nl

Abstract

Human visceral leishmaniasis (VL) vaccines are currently under development and there is a need to understand their potential impact on population wide VL incidence. We implement four characteristics from different human VL vaccine candidates into two published VL transmission model variants to estimate the potential impact of these vaccine characteristics on population-wide anthroponotic VL incidence on the Indian subcontinent (ISC). The vaccines that are simulated in this study 1) reduce the infectiousness of infected individuals towards sand flies, 2) reduce risk of developing symptoms after infection, 3) reduce the risk of developing post-kala-azar dermal leishmaniasis (PKDL), or 4) lead to the development of transient immunity. We also compare and combine a vaccine strategy with current interventions to identify their potential role in elimination of VL as a public health problem. We show that the first two simulated vaccine characteristics can greatly reduce VL incidence. For these vaccines, an approximate 60% vaccine efficacy would lead to achieving the ISC elimination target (<1 VL case per 10,000 population per year) within 10 years' time in a moderately endemic setting when vaccinating 100% of the population. Vaccinating VL cases to prevent the development of PKDL is a promising tool to sustain the low incidence elimination target after regular interventions are halted. Vaccines triggering the development of transient immunity protecting against infection lead to the biggest reduction in VL incidence, but booster doses are required to achieve perduring impact. Even though vaccines are not yet available for implementation, their development should be pursued as their potential impact on transmission can be substantial, both in decreasing incidence at the population level as well as in sustaining the ISC elimination target when other interventions are halted.

Author summary

Vaccines for human visceral leishmaniasis (VL) are currently under development. In this study, we simulate VL transmission dynamics using mathematical models to explore the potential impact of vaccines on population-wide incidence. We show that some vaccines have high potential to reduce VL incidence, namely those that reduce the infectiousness of infected individuals to sand flies and those that reduce the chance of developing symptoms once infected. The effect of vaccines that lead to protection from infection is potentially the greatest, but depending on the duration of immunity, individuals would require booster doses to guarantee lifelong impact. Vaccines that prevent the development of post-kala-azar dermal leishmaniasis are a promising tool to sustain low VL incidence and prevent recrudescence of infection when regular interventions are halted. Our results strongly support the continued development of VL vaccines, as their potential impact on population incidence can be substantial.

Introduction

Visceral leishmaniasis (VL), also known as kala-azar, is a vector-borne neglected tropical disease. Infection occurs after successful transmission of the *Leishmania* protozoa through the bite of an infected female sand fly [1]. Most infected humans remain asymptomatic, and only a small proportion of about 1—10% develop clinical symptoms, resulting in death when left untreated [2,3]. Between 5% and 20% of treated VL cases develop a long-lasting skin condition known as post-kala-azar dermal leishmaniasis (PKDL). Recent studies have identified that individuals with PKDL are equally infectious towards sand flies as VL cases, making them an important reservoir of infection [4,5]. However, the contribution of asymptomatic individuals to transmission has not yet been defined [4,6]. After infection, a period of immunity follows, of which the duration remains debated.

Currently around 33,000—66,000 individuals develop symptomatic VL each year, mainly on the Indian subcontinent (ISC), Eastern Africa, the Mediterranean region, and Brazil, affecting the poorest of the poor [7,8]. The World Health Organization (WHO) and affected countries target for ‘elimination of VL as a public health problem by 2020’ on the ISC, where VL is considered to be solely anthroponotic. This target is defined as maintaining less than 1 VL case per 10,000 individuals per year at district level in Nepal, at subdistrict/block level in India, and at upazila level in Bangladesh [9]. In the rest of the world (e.g Africa, Europe, Brazil), where VL can also be zoonotic with the main reservoir of infection in dogs, the target is 100% detection and treatment of symptomatic cases [10]. Current strategies consist of diagnosis and treatment of VL cases, and vector control.

Vaccines already play an important role in the control of canine leishmaniasis, at the

individual level they reduce the development of symptoms, reduce the parasite load in the blood, and reduce the risk of death [11–13]. These vaccines have also proven to be effective at the population level by reducing *Leishmania* transmission, resulting in lower incidence in both dogs and humans [14,15]. The development of human VL vaccines has been on-going for decades and there are different vaccine candidates currently in trial, but none are yet available for implementation [16,17]. The promising results from experimental human VL vaccine trials, and by the practice of “leishmanization”, in which a healthy individual is artificially exposed to tissue scrapings derived from a cutaneous leishmaniasis patient, leading to disease prevention [6,16,18–20], provide strong evidence for the scientific feasibility of an effective vaccine against human VL. Should an effective vaccine become available, it has been estimated to be cost-effective when used at large scale and in addition to ongoing diagnosis and treatment, without even accounting for its impact on transmission [19].

Mathematical transmission models are useful tools to gain insight into the effect of current and future interventions on VL incidence and the underlying transmission dynamics. Previous modelling studies that focused on VL transmission on the ISC presented two model variants; one in which only VL and PKDL cases contribute to transmission, and another in which also asymptomatic individuals contribute to transmission (~1% relative to VL cases). The models estimated that in most situations on the ISC, the target is likely to be met with current strategies but in high endemic settings and at a lower geographical scale, additional efforts are required. They also highlighted the risk of recrudescence of infection after achieving the low incidence target, when halting interventions. This is mainly due to individuals with PKDL and/or asymptomatic infection. Therefore, the studies emphasized the need for further research on the potential impact of preventive VL and PKDL strategies as a

tool in reaching and sustaining VL elimination as a public health problem on the ISC [5,21,22]. Other studies stressed that 100% detection and treatment of cases in the rest of the world remains challenging and that prevention could be much more effective than case detection and treatment [23].

In this study, we implement multiple characteristics of potential human VL vaccines using the two variants of a deterministic VL transmission model [21] to estimate the potential impact of these vaccine characteristics on VL incidence and transmission dynamics during and after the achievement of the current elimination target. The vaccines that are simulated in this study 1) reduce the infectiousness of infected individuals towards the sand fly, 2) reduce the risk of developing symptoms after infection, 3) reduce the risk of development of PKDL, or 4) lead to the development of transient immunity to infection [24–26]. We also compare and combine vaccine characteristics with current interventions to identify which vaccines could be most impactful in fighting this neglected tropical disease.

Methods

Overview of VL vaccine candidates and characteristics

Currently there are various VL vaccine candidates under study [27]: LEISH-F3+GLA-SE [28,29], and ChAd63-KH (ISRCTN07766359) [30] are currently in clinical development; Ad5-A2/rA2 Prime / Boost [31], genetically modified live attenuated whole parasites [25,26,32], and a LmCen^{-/-} vaccine [33] are being developed for the clinic [34].

These vaccines have different physical and immunological properties, and could be used in either prophylactic or therapeutic settings, but their impact following infected sand fly bite in

humans has yet to be evaluated. Table 1 summarizes different potential vaccine outcome measures (herein called characteristics) that were selected for simulation in this study. Vaccine characteristic 1 is separated into 1a) asymptomatic individuals and 1b) all infected individuals, because it is suggested that only individuals with asymptomatic infection may be affected by the vaccine and that once an individual develops symptoms there are no differences in infectiveness (1a). However, since this is not yet well established, we also include the option where all infected individuals become less infective, as a result of the vaccine (1b).

Table 1. Human VL vaccine characteristics.

Number	Vaccine characteristic
1a	Reduced infectivity of asymptomatic individuals
1b	Reduced infectivity of all infected individuals
2	Reduced risk of developing symptoms
3	Reduced risk of developing PKDL
4	Development of transient immunity protecting against infection

Transmission models and simulation of existing interventions

Fig 1 illustrates the basic structure of the VL transmission model, which is a deterministic age-structured model. There are two model variants, that only differ based on assumptions about where the main reservoir of infection lies; namely, solely in symptomatic individuals (VL and PKDL), or mainly in asymptomatic individuals [21,35,36]. The models were parameterized with age-structured data on approximately 21,000 individuals included in the KalaNet bednet trial in India and Nepal [37] and have undergone geographical cross-validation against data on >5000 VL cases from 8 endemic districts in Bihar collected by CARE India [38] (see [36] for full model code and descriptions, and sensitivity analyses).

Recent outcomes from xenodiagnosis studies have been incorporated, indicating that those with PKDL are on average nearly as infectious as those with VL (0.9:1.0) [4,5].

Interventions of which the effects have previously been modelled are vector control through indoor-residual spraying of insecticide (IRS) and active case detection (ACD). The guidelines, as developed by WHO, recommend a 5-year attack phase (intense IRS and ACD) followed by 5 years of consolidation phase (IRS and intense ACD). In our models, IRS leads to a decrease in sand fly density and ACD shortens the duration of the symptomatic untreated stage.

Implementation and simulation of four vaccine characteristics

Vaccine characteristic 1 is simulated by a reduction in infectiousness of infectious states towards the sand fly. For vaccine characteristics 2 and 3, the respective flow towards clinical VL and PKDL is reduced. With vaccine characteristic 4, we selected 100% development of transient immunity after having received the vaccine and experimented with vaccinating 100% and 50% of the population. The duration of immunity after vaccination is assumed to be 2 years, which is similar to the assumed duration of immunity after natural infection in our model of which sensitivity analyses are presented in previous work [36].

For the simulations of vaccine characteristics, we assume that they apply to everyone involved, i.e. all ages and sexes. No specific target populations are simulated, besides for vaccine characteristic 3, which is only administered to those that have developed VL. For vaccine characteristics 1-3, we assume an arbitrary 50% reduction of the infectiousness as well as a 50% reduction of the proportions of individuals that develop VL and PKDL, all in combination with a 100% vaccination coverage. We also calculate the percentage of vaccine

effectiveness required to achieve the VL elimination target incidence of 1/10,000/year within 10 years of starting the intervention, which could aid in defining a vaccine target product profile (TPP). We assume that the vaccine characteristics are in place constantly from the start of the intervention, except for vaccine characteristic 4, where we experiment with simulating a single vaccination round and repeated yearly vaccination rounds. For all four vaccine characteristics, we separately simulate and compare their impact on VL incidence over time, even though it is likely that one vaccine will possess multiple characteristics. The cumulative effects of some vaccine characteristics are simulated indirectly, as reducing the development of VL will lead to a decrease in the overall development of PKDL. Previous work has shown that when current existing interventions have led to the 1/10,000/year target, there are many susceptible individuals and the infection pressure comes mainly from PKDL cases (when assuming the infection pressure originates from symptomatic individuals only) [21]. To address this, we also combine vaccine characteristic 3, vaccination of VL cases to prevent the development of PKDL, with the current interventions recommended by WHO.

Fig 1. Schematic presentation of the model variant in which asymptomatic individuals contribute to transmission, with numbers related to different types of vaccine characteristics that are implemented in the models. In the alternative model variant, asymptomatic individuals are assumed not to be infectious towards to sand fly, with infection pressure only coming from symptomatic individuals with VL (with and without treatment) and PKDL. Once a susceptible individual is infected by an infectious sand fly, they become early asymptomatic for an average duration of about 200 days, which is followed by the late asymptomatic stage (average duration of 69 days). The average infectivity of both asymptomatic stages together is 0 in the model in which they do not contribute and ~1.5% relative to VL in the model in which they contribute to transmission. 1.4% of late asymptomatic individuals develops VL, and without active case detection, the duration between onset of symptoms and start of treatment lasts on average 40 days, followed by 1-day treatment 1 and potentially 28-day treatment 2 or death if left untreated. The average duration of the putatively recovered stage is 21 months and 5% of these individuals develop PKDL which lasts 5 years on average. The infectivity of PKDL is 90%, relative to VL. The rest recovers to the early recovered stage (average duration of 74 days), followed by the late recovered stage (average duration of 2 years), which can be interpreted as the duration of immunity. The numbers in the red

boxes relate to the numbers in the first column of Table 1 and represent the following vaccine characteristics; 1a) early and late asymptomatic individuals become half as infectious, 1b) all infection states become half as infectious, 2) vaccinated individuals are 50% less likely to develop symptoms, 3) vaccinated individuals are 50% less likely to develop PKDL, and 4) vaccinated individuals develop transient immunity against infection.

Results

The impact of each of the four vaccine characteristics on VL incidence is illustrated in Fig 2. A vaccine that reduces infectivity of asymptomatic individuals by 50% (1a) leads to achievement of the target of less than 1 VL case per 10,000 population per year in about 11 years. When all infected individuals have a reduced infectiousness of 50% (1b), the decline is steeper, achieving elimination in around 4 years if asymptomatics are the main reservoir of infection and 11 years when infection is only coming from those with VL and PKDL. Halving the chance of developing symptoms (2) also has a considerable impact on transmission, especially if only symptomatic individuals are infective after which elimination takes about 10 years. However, if most infection pressure arises from asymptomatic individuals, the impact of halving the development of symptoms will lead to achieving the elimination target only after about 19 years, when used as a stand-alone tool. A 50% reduction in the development of PKDL (3), after which not 5% (default) but only 2.5% of past VL cases develop PKDL, has the smallest impact on transmission. As expected with this characteristic, the relatively larger impact is seen when only those with VL and PKDL contribute to transmission, and thus when PKDL plays a more prominent role in the transmission dynamics. Of all vaccine characteristics, the development of immunity that protects against infection (as seen in late recovered cases) of the population causes the most rapid decrease in incidence (4), since the pool of susceptible individuals is completely removed at once (with the assumption of 100% coverage as used in the model). We

230 additionally explored the effect of vaccinating half the population and repeating this yearly
231 for 5 years in a row (5 x 50%), showing that regular vaccinations are required to sustain the
232 impact and move towards the low incidence elimination target.

233
234
235 **Fig 2. The impact of different vaccine characteristics on VL incidence using model variants with and without**
236 **asymptomatics contributing to transmission in a setting with a pre-control endemicity of 5/10,000/year.** Vaccine
237 characteristics are in place continuously from year 0 onwards, unless for vaccine characteristic 4, which is administered once
238 (1 x 100%), or yearly for five years in a row (5 x 50%). The different vaccine characteristics that are also explained in Table
239 1 and illustrated in Fig 1, are; 1a) early and late asymptomatic individuals become half as infectious, 1b) all infection states
240 become half as infectious, 2) vaccinated individuals are 50% less likely to develop symptoms, 3) vaccinated individuals are
241 50% less likely to develop PKDL, and 4) vaccinated individuals become immediately immune. The black dashed line
242 represents the WHO elimination target of 1/10,000/year. The oscillations in VL incidence are a result of seasonality in the
243 sand fly density.

244
245 The minimum vaccine effect required for each vaccine characteristic to achieve the VL
246 elimination target incidence of 1/10,000/year within 10 years of starting the intervention is
247 presented in Table 2. The vaccine characteristics that impact the development of VL and
248 PKDL (2 and 3) obviously have a bigger impact in the model in which only VL and PKDL
249 contribute to transmission.

Table 2. Minimum required effect of the vaccine characteristics to reach a VL elimination target incidence of 1/10,000/year within 10 years' time after starting the intervention, when vaccinating 100% of the population in a setting with a 5/10,000/year pre-control incidence.

Vaccine characteristic	Model variant	
	Only VL and PKDL contribute to transmission	Asymptomatics are main contributors to transmission
1a) required reduction in infectivity of asymptomatic individuals	N/A	35%
1b) required reduction in infectivity of all infected individuals	60%	37%
2) required reduction in the development of symptoms	56%	68%
3) required amount of time to reach the elimination target when preventing the development of PKDL completely	11 years	>20 years
4) required minimum number of rounds when vaccinating 50% of the susceptible individuals yearly with 100% vaccine efficacy	14 rounds	5 rounds

Vaccine characteristic 3, after which vaccinated individuals are less likely to develop PKDL, displayed the least impact when used as a stand-alone tool. Fig 3 shows the impact on VL incidence of a decrease in the development of PKDL of 50% and 100%, combined with the current interventions for a setting with a pre-control endemicity level of 5/10,000/year. The red line represents the default scenario in which the current interventions (active case detection and vector control) are in place during the WHO attack phase (year 0—5) and the WHO consolidation phase (year 5—10), without the presence of a vaccine. Further details on the impact of current interventions on VL incidence on the ISC as predicted by these models can be found in Le Rutte *et al.*, 2018 [21]. After halting all interventions at year 10, the situation will slowly return to the pre-control equilibrium of 5/10,000/year, because of the remaining VL incidence in year 10 in all scenarios. In the two scenarios with the PKDL vaccine (green and blue lines) a new, much lower, equilibrium will be reached after regular interventions are halted. For the vaccine with a 50% efficacy (50% decrease in PKDL

development of vaccinated VL cases) the target of 1/10,000/year will be reached as simulated by the model in which only VL and PKDL contribute to transmission. When assuming an effect of 100% protection from developing PKDL, this model suggests that using only vaccine 3 could keep the incidence below 1/10,000/year, after all regular interventions have brought incidence down and are halted. However, in settings with a higher pre-control endemicity of 10/10,000/year, only the vaccine with 100% protection against development of PKDL will lead to the elimination target of VL after 15-20 years depending on the start year of the PKDL vaccine.

Fig 3. Strategies of combining vaccine effect 3 with the WHO attack and consolidation phase for a setting with a pre-control endemicity level of 5/10,000/year. Top panels: vaccine effect with 100% protection against the development of PKDL, bottom panels: vaccine effect with 50% protection against the development of PKDL. The default strategy is visualized with the red line (top and bottom row identical), in which 5 years of attack phase are followed by 5 years of consolidation phase, after which interventions are halted in year 10. For the green line, the PKDL vaccine is introduced during the consolidation phase (year 5), which continues after the consolidation phase has ended at year 10. For the blue line, the PKDL vaccine is already introduced at the start of the attack phase (year 0), continues during the consolidation phase and is continued when regular interventions are halted in year 10. Left figures show the simulations for the model variant where solely symptomatic individuals contribute to transmission, whereas for the right figures asymptomatic individuals constitute the main reservoir of infection. The black dashed line represents the WHO VL incidence target of 1/10,000/year. The oscillations in VL incidence are a result of seasonality in the sand fly density.

Discussion

In this study, we present for the first time the potential impact of VL vaccines on transmission dynamics and population incidence on the Indian subcontinent (ISC). This impact looks very promising. We found that all simulated vaccine characteristics show potential in reducing population VL incidence, particularly those that reduce the infected individual's infectiousness or reduce the chance of developing symptoms once infected. For

these vaccines, an approximate 60% vaccine efficacy would lead to achieving the ISC elimination target (<1 VL case per 10,000 population per year) within 10 years' time in a moderately endemic setting, assuming that the entire population is vaccinated and only VL and PKDL cases contribute to transmission. For the model variant in which asymptomatics are the main contributors to transmission, much lower vaccine efficacies of around 37% would be required when reducing the infectiousness; however, for the required reduction in the development of symptoms, a vaccine efficacy of nearly 70% was estimated. The vaccine that leads to immunity akin to that of late recovered cases shows the highest impact, but individuals would require regular booster vaccines to achieve and sustain the low incidence elimination target. Vaccinating VL cases to prevent the development of PKDL shows to be a promising tool to sustain the elimination target once reached, and prevent recrudescence of infection when regular interventions are halted. Those findings are of great importance in providing a factual base to the ongoing effort aimed at establishing a TPP for a VL vaccine.

A limitation to our study is the fact that we simulated vaccine characteristics rather simplistically by instantaneously altering the transition rates and applying this simultaneously to all individuals in the population. Ideally, vaccinated individuals should move to different, additional, compartments in the model, where they experience a different history of infection. In such a model, vaccinated and unvaccinated individuals would be living beside each other, both influencing the transmission dynamics differently, although the outcomes would likely only differ quantitatively with ours. Another limitation of our study is that we only present the results for a setting with a pre-control endemicity of 5 VL cases per 10,000 population per year, which we considered representative for endemic situations where vaccines would be most useful. In settings with a lower pre-control endemicity the elimination target would be achieved earlier; in settings with a higher pre-control endemicity, the vaccine characteristics

would require a higher efficacy to achieve the same effect on VL incidence in the same amount of time.

We further decided to simulate the vaccine characteristics separately, while in reality most vaccines are expected to possess multiple characteristics. For example lowering the parasite load will likely lead to both decreased infectiousness as well as reduced development of symptoms, as is also seen in canine VL vaccines [12,13]. However, by combining them it would be less clear to what extent different characteristics would drive the total impact of a vaccine. For the vaccine that causes vaccinated individuals to develop transient immunity against infection, it is important to note that the impact on VL incidence, as well as the required number of booster vaccines, highly depends on the duration of acquired immunity, which was assumed to be two years on average in our models similar to what we used in previous work [36]. The longer the duration of acquired immunity, the bigger the impact on VL incidence and the lower the frequency of required booster vaccines. We also assume that for all vaccine characteristics the efficacy is 50%. Even though this is a generalization and in reality it is likely different for each characteristic, this approach allows us to compare the impact of the different vaccine characteristics. In this study we simulate transmission between humans and sand flies, which is currently considered to reflect the transmission dynamics of VL on the ISC. However, would a considerable contribution to transmission come from an animal reservoir, vertical transmission as seen in dogs, and/or the presence of those with HIV-VL co-infection, the potential impact of vaccines could increase [13,39–41].

A typical aspect of the deterministic model that we use is that all durations of states are exponentially distributed, which often does not reflect the actual distributions of durations as found in nature. The slow recrudescence of infection between year 10 and 20 is another

phenomenon of the deterministic model, where prevalences can never become completely zero, but in reality the disease will either die out or come back, and if it comes back, most likely it will progress somewhat faster. Around the elimination target when numbers of infected cases become very low, the role of chance increases and a stochastic transmission model would be required to analyse the risks of recrudescence or the chance of achieving (local) elimination of transmission.

We acknowledge that some of the assumptions chosen for the simulation are not fully reflective of the reality of implemented immunization programs. Firstly, our choice of 100% coverage certainly is an overestimation of what can be realistically achieved. For example, coverage for the 1st dose of measles-containing vaccines was on average 73% in the AFRO region, and the human papilloma virus vaccination had an average coverage of 88% when pooling regions and income levels [42,43]. However, this assumption allowed us to evaluate the maximum impact and to do such an evaluation independently from the constraints of delivery strategy. Also, having chosen another coverage level would not alter our outcomes when comparing the impact of the different characteristics. With a lower coverage, the durations until elimination as a public health problem would be longer and the minimum required efficacies would be higher. Secondly, and in particular at the start of vaccination programs, a catch up campaign is usually implemented to quickly reduce the susceptible population, focusing on the population that is at highest risk (i.e. for leishmaniasis, children and young adults or migrant workers [44]). Such programmatic design considerations are not considered in the current model and will need to be investigated with more complex individual-based transmission models. Lastly, 5-year protection is most likely going to be the minimum requirement to allow for a widespread roll-out in routine immunization. Shorter durations requiring a very frequent administration of booster doses might prove

programmatically and financially unsustainable. Nevertheless, from an impact assessment standpoint the results generated with the more conservative assumptions of the current model have clear significance for understanding the relative importance of different vaccine characteristics.

Vaccines have proven to be vital tools in the control and prevention of diseases [45,46]. This study reveals that a VL vaccine strategy could also prove an important tool in the fight against this neglected tropical disease. We focussed on the anthroponotic transmission dynamics of VL on the Indian subcontinent, but also in the rest of the world VL vaccines are likely to surpass their impact at the patient level by reducing the infection pressure, positively impacting the estimated 6 million people at risk of VL globally [47].

In conclusion, even though VL vaccines are not yet available for implementation, our results strongly support their continued development, given the potentially substantive impact on transmission, decreasing incidence at the population level, and sustaining the low incidence elimination target on the ISC when other interventions are relaxed. More details of the impact of different vaccines characteristics on the history of infection are awaited to further our understanding and modelling of the impact of VL vaccines on VL transmission dynamics and disease incidence.

Acknowledgements

We are very grateful for having had the opportunity to learn from the insights and tremendous expertise of Professor Farrokh Modabber while discussing this work. We would also like to thank the organisers and attendees of the VL vaccine expert meeting in Rockville,

USA, in September 2015 at the National Institute of Allergy and Infectious Diseases, where the idea for this study sparked.

References

1. WHO SEARO. WHO | Leishmaniasis fact sheet. 2019 [cited 1 Oct 2019]. Available: http://www.searo.who.int/entity/vector_borne_tropical_diseases/data/FactSheetVL.pdf
2. Ostyn B, Gidwani K, Khanal B, Picado A, Chappuis F, Singh SP, et al. Incidence of symptomatic and asymptomatic *Leishmania donovani* infections in High-Endemic foci in India and Nepal: A prospective study. *PLoS Negl Trop Dis*. 2011;5: 1–7. doi:10.1371/journal.pntd.0001284
3. Le Rutte EA, Coffeng LE, Bontje DM, Hasker EC, Ruiz Postigo JA, Argaw D, et al. Feasibility of eliminating visceral leishmaniasis from the Indian subcontinent: Explorations with a set of deterministic age-structured transmission models Quantitative analysis of strategies to achieve the 2020 goals for neglected tropical diseases: *Wher. Parasites and Vectors*. 2016;9. doi:10.1186/s13071-016-1292-0
4. Mondal D, Bern C, Ghosh D, Rashid M, Molina R, Chowdhury R, et al. Quantifying the Infectiousness of Post-Kala-Azar Dermal Leishmaniasis Toward Sand Flies. *Clin Infect Dis*. 2019;69: 251–258. doi:10.1093/cid/ciy891
5. Le Rutte EA, Zijlstra EE, de Vlas SJ. Post-Kala-Azar Dermal Leishmaniasis as a Reservoir for Visceral Leishmaniasis Transmission. *Trends Parasitol*. 2019;35: 590–592. doi:10.1016/j.pt.2019.06.007
6. Nagill R, Kaur S. Vaccine candidates for leishmaniasis: A review. *Int Immunopharmacol*. 2011. doi:10.1016/j.intimp.2011.05.008
7. World Health Organization. Leishmaniasis Fact sheet. 2018 [cited 20 Dec 2018]. Available: <http://www.who.int/en/news-room/fact-sheets/detail/leishmaniasis>

- 420 8. James SL, Abate D, Abate KH, Abay SM, Abbafati C, Abbasi N, et al. Global, regional, and
421 national incidence, prevalence, and years lived with disability for 354 diseases and injuries for
422 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of
423 Disease Study 2017. *Lancet*. 2018;392: 1789–1858. doi:10.1016/S0140-6736(18)32279-7
- 424 9. World Health Organization Regional Office for South-East Asia. Process of validation of
425 elimination of kala-azar SEARO. 2016. Available:
426 [https://www.who.int/leishmaniasis/resources/Process_of_validation_of_VL_elimination_SEA](https://www.who.int/leishmaniasis/resources/Process_of_validation_of_VL_elimination_SEA_CD_321.pdf?ua=1&ua=1)
427 [_CD_321.pdf?ua=1&ua=1](https://www.who.int/leishmaniasis/resources/Process_of_validation_of_VL_elimination_SEA_CD_321.pdf?ua=1&ua=1)
- 428 10. World Health Organisation. Integrating Neglected Tropical Diseases into global health and
429 development. 2017.
- 430 11. Trigo J, Abbehusen M, Netto EM, Nakatani M, Pedral- G, Jesus RS De, et al. Treatment of
431 canine visceral leishmaniasis by the vaccine Leish-111f+MPL-SE Joelma. 2011;28: 3333–
432 3340. doi:10.1016/j.vaccine.2010.02.089.Treatment
- 433 12. Palatnik-de-Sousa CB. Vaccines for Canine Leishmaniasis. *Front Immunol*. 2012;3.
434 doi:10.3389/fimmu.2012.00069
- 435 13. Toepp A, Larson M, Wilson G, Grinnage-Pulley T, Bennett C, Leal-Lima A, et al.
436 Randomized, controlled, double-blinded field trial to assess *Leishmania* vaccine effectiveness
437 as immunotherapy for canine leishmaniosis. *Vaccine*. 2018;36: 6433–6441.
438 doi:10.1016/j.vaccine.2018.08.087
- 439 14. Kumar R, Engwerda C. Vaccines to prevent leishmaniasis. *Clin Transl Immunol*. 2014;3: e13.
440 doi:10.1038/cti.2014.4
- 441 15. Jain K, Jain NK. Vaccines for visceral leishmaniasis: A review. *J Immunol Methods*.
442 2015;422: 1–12. doi:10.1016/j.jim.2015.03.017
- 443 16. Jain K, Jain NK. Vaccines for visceral leishmaniasis: A review. *J Immunol Methods*. 2015.
444 doi:10.1016/j.jim.2015.03.017

- 445 17. Modabber F. Leishmaniasis vaccines: past, present and future. *Int J Antimicrob Agents*.
446 2010;36: S58–S61. doi:10.1016/j.ijantimicag.2010.06.024
- 447 18. Chappuis F, Sundar S, Hailu A, Ghalib H, Rijal S, Peeling RW, et al. Visceral leishmaniasis:
448 what are the needs for diagnosis, treatment and control? *Nat Rev Microbiol*. 2007/10/17.
449 2007;5: 873–882. doi:10.1038/nrmicro1748
- 450 19. Lee BY, Bacon KM, Shah M, Kitchen SB, Connor DL, Slayton RB. The Economic Value of a
451 Visceral Leishmaniasis Vaccine in Bihar State, India. *Am J Trop Med Hyg*. 2012.
452 doi:10.4269/ajtmh.2012.10-0415
- 453 20. Khamesipour A, Dowlati Y, Asilian A, Hashemi-Fesharki R, Javadi A, Noazin S, et al.
454 Leishmanization: Use of an old method for evaluation of candidate vaccines against
455 leishmaniasis. *Vaccine*. 2005. doi:10.1016/j.vaccine.2005.02.015
- 456 21. Le Rutte EA, Chapman LAC, Coffeng LE, Ruiz-Postigo JA, Olliaro PL, Adams ER, et al.
457 Policy Recommendations From Transmission Modeling for the Elimination of Visceral
458 Leishmaniasis in the Indian Subcontinent. *Clin Infect Dis*. 2018;66: S301–S308.
459 doi:10.1093/cid/ciy007
- 460 22. Sundar S, Singh OP, Chakravarty J. Visceral leishmaniasis elimination targets in India,
461 strategies for preventing resurgence. *Expert Rev Anti Infect Ther*. 2018;16: 805–812.
462 doi:10.1080/14787210.2018.1532790
- 463 23. Boelaert M, Criel B, Leeuwenburg J, Van Damme W, Le Ray D, Van der Stuyft P. Visceral
464 leishmaniasis control: a public health perspective. *Trans R Soc Trop Med Hyg*. 2000;94: 465–
465 471. doi:10.1016/S0035-9203(00)90055-5
- 466 24. Oliveira F, Rowton E, Aslan H, Gomes R, Castrovinci PA, Alvarenga PH, et al. A sand fly
467 salivary protein vaccine shows efficacy against vector-transmitted cutaneous leishmaniasis in
468 nonhuman primates. *Sci Transl Med*. 2015;7: 290ra90. doi:10.1126/scitranslmed.aaa3043
- 469 25. Zahedifard F, Gholami E, Taheri T, Taslimi Y, Doustdari F, Seyed N, et al. Enhanced

- 470 Protective Efficacy of Nonpathogenic Recombinant *Leishmania tarentolae* Expressing
 471 Cysteine Proteinases Combined with a Sand Fly Salivary Antigen. McMahon-Pratt D, editor.
 472 PLoS Negl Trop Dis. 2014;8: e2751. doi:10.1371/journal.pntd.0002751
- 473 26. Dey R, Natarajan G, Bhattacharya P, Cummings H, Dagur PK, Terrazas C, et al.
 474 Characterization of Cross-Protection by Genetically Modified Live-Attenuated *Leishmania*
 475 *donovani* Parasites against *Leishmania mexicana*. J Immunol. 2014;193: 3513–3527.
 476 doi:10.4049/jimmunol.1303145
- 477 27. Mo AX, Pesce J, Fenton Hall B. Meeting report: Visceral leishmaniasis control and
 478 elimination: Is there a role for vaccines in achieving regional and global goals? Am J Trop
 479 Med Hyg. 2016;95: 514–521. doi:10.4269/ajtmh.16-0184
- 480 28. Coler RN, Duthie MS, Hofmeyer KA, Guderian J, Jayashankar L, Vergara J, et al. From
 481 mouse to man: safety, immunogenicity and efficacy of a candidate leishmaniasis vaccine
 482 LEISH-F3+GLA-SE. Clin Transl Immunol. 2015. doi:10.1038/cti.2015.6
- 483 29. Duthie MS, Pereira L, Favila M, Hofmeyer KA, Reed SJ, Metangmo S, et al. A defined
 484 subunit vaccine that protects against vector-borne visceral leishmaniasis. NPJ vaccines.
 485 2017;2: 23. doi:10.1038/s41541-017-0025-5
- 486 30. Osman M, Mistry A, Keding A, Gabe R, Cook E, Forrester S, et al. A third generation vaccine
 487 for human visceral leishmaniasis and post kala azar dermal leishmaniasis: First-in-human trial
 488 of ChAd63-KH. McDowell MA, editor. PLoS Negl Trop Dis. 2017;11: e0005527.
 489 doi:10.1371/journal.pntd.0005527
- 490 31. Grimaldi G, Teva A, Porrozzi R, Pinto M a, Marchevsky RS, Rocha MGL, et al. Clinical and
 491 Parasitological Protection in a *Leishmania infantum*-Macaque Model Vaccinated with
 492 Adenovirus and the Recombinant A2 Antigen. Nakhasi HL, editor. PLoS Negl Trop Dis.
 493 2014;8: e2853. doi:10.1371/journal.pntd.0002853
- 494 32. Ismail N, Kaul A, Bhattacharya P, Gannavaram S, Nakhasi HL. Immunization with Live
 495 Attenuated *Leishmania donovani* Centrin–/– Parasites Is Efficacious in Asymptomatic

496 Infection. Front Immunol. 2017;8. doi:10.3389/fimmu.2017.01788

497 33. Ismail N, Karmakar S, Bhattacharya P, Dey R, Nakhasi HL. Immunization with Leishmania
498 major centrin knock-out (LmCen^{-/-}) parasites induces skin resident memory T cells that plays
499 a role in protection against wild type infection (LmWT). J Immunol. 2019;202: 196.29 LP-
500 196.29. Available: http://www.jimmunol.org/content/202/1_Supplement/196.29.abstract

501 34. Global Health Innovative Technology Fund. Live attenuated prophylactic vaccine for
502 leishmaniasis - Investment Details. 2018 [cited 11 Nov 2019]. Available:
503 <https://www.ghitfund.org/investment/portfoliodetail/detail/135>

504 35. Le Rutte EA, Coffeng LE, Bontje DM, Hasker EC, Ruiz Postigo JA, Argaw D, et al.
505 Feasibility of eliminating visceral leishmaniasis from the Indian subcontinent: explorations
506 with a set of deterministic age-structured transmission models. Parasit Vectors. 2016;9: 24.
507 doi:10.1186/s13071-016-1292-0

508 36. Le Rutte EA, Chapman LAC, Coffeng LE, Jervis S, Hasker EC, Dwivedi S, et al. Elimination
509 of visceral leishmaniasis in the Indian subcontinent: a comparison of predictions from three
510 transmission models. Epidemics. 2017;18: 67–80. doi:10.1016/j.epidem.2017.01.002

511 37. Picado A, Singh SP, Rijal S, Sundar S, Ostyn B, Chappuis F, et al. Longlasting insecticidal
512 nets for prevention of Leishmania donovani infection in India and Nepal: paired cluster
513 randomised trial. BMJ. 2010;341: c6760. doi:10.1136/bmj.c6760

514 38. Jervis S, Chapman LAC, Dwivedi S, Karthick M, Das A, Le Rutte EA, et al. Variations in
515 visceral leishmaniasis burden, mortality and the pathway to care within Bihar, India. Parasit
516 Vectors. 2017;10: 601. doi:10.1186/s13071-017-2530-9

517 39. Burza S, Mahajan R, Sanz MG, Sunyoto T, Kumar R, Mitra G, et al. HIV and Visceral
518 Leishmaniasis Coinfection in Bihar, India: An Underrecognized and Underdiagnosed Threat
519 Against Elimination. Clin Infect Dis. 2014;59: 552–555. doi:10.1093/cid/ciu333

520 40. Singh N, Mishra J, Singh R, Singh S. Animal reservoirs of visceral leishmaniasis in India. J

Parasitol. 2012/07/07. 2013;99: 64–67. doi:10.1645/ge-3085.1

41. Meinecke CK, Schottelius J, Oskam L, Fleischer B. Congenital Transmission of Visceral Leishmaniasis (Kala Azar) From an Asymptomatic Mother to Her Child. *Pediatrics*. 1999;104: e65–e65. doi:10.1542/peds.104.5.e65

42. Bruni L, Diaz M, Barrionuevo-Rosas L, Herrero R, Bray F, Bosch FX, et al. Global estimates of human papillomavirus vaccination coverage by region and income level: a pooled analysis. *Lancet Glob Heal*. 2016;4: e453–e463. doi:10.1016/S2214-109X(16)30099-7


43. WHO/UNICEF. Coverage estimates for AFRO region. 2018. Available: www.who.int/immunization/monitoring_surveillance/en/

44. Leta S, Dao THT, Mesele F, Alemayehu G. Visceral Leishmaniasis in Ethiopia: An Evolving Disease. Ghedin E, editor. *PLoS Negl Trop Dis*. 2014;8: e3131. doi:10.1371/journal.pntd.0003131

45. Bulletin of the World Health Organization. Vaccination greatly reduces disease, disability, death and inequity worldwide. In: 2008.

46. Muller CP, Kremer JR, Best JM, Dourado I, Triki H, Reef S. Reducing global disease burden of measles and rubella: Report of the WHO Steering Committee on research related to measles and rubella vaccines and vaccination, 2005. *Vaccine*. 2007;25: 1–9. doi:10.1016/j.vaccine.2006.07.039

47. WHO. Leishmaniasis in high-burden countries: an epidemiological update based on data reported in 2014. *Wkly Epidemiol Rec*. 2016; 285–296. doi:10.1186/1750-9378-2-15.Voir



Model predictions

- Only VL and PKDL contribute to transmission
- Asymptomatics are main contributors to transmission

